

# *Systematic under- and overestimation of GHG reductions in renewable biomass systems*

## **Climatic Change**

An Interdisciplinary,  
International Journal Devoted  
to the Description, Causes and  
Implications of Climatic Change

ISSN 0165-0009

Volume 104

Number 2

Climatic Change (2010)

104:415-422

DOI 10.1007/

s10584-010-9984-5

## Climatic Change

An Interdisciplinary, International Journal Devoted to the  
Description, Causes and Implications of Climatic Change

*Editors:* **MICHAEL OPPENHEIMER**  
**GARY YOHE**

Volume 104 – No. 2 – January II 2011

Including **CLIMATIC CHANGE LETTERS**  
*Editor:* **Michael Oppenheimer**



 Springer

ISSN 0165-0009

 Springer

**Your article is protected by copyright and all rights are held exclusively by Springer Science+Business Media B.V.. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your work, please use the accepted author's version for posting to your own website or your institution's repository. You may further deposit the accepted author's version on a funder's repository at a funder's request, provided it is not made publicly available until 12 months after publication.**

# Systematic under- and overestimation of GHG reductions in renewable biomass systems

## A Letter

Thea L. Whitman · C. Johannes Lehmann

Received: 16 July 2010 / Accepted: 29 October 2010 / Published online: 26 November 2010  
© Springer Science+Business Media B.V. 2010

**Abstract** This paper identifies a critical systematic error in greenhouse gas accounting in renewable biomass systems. While CO<sub>2</sub> emissions from renewable biomass energy systems are generally considered to have a net impact of 0, no similar adjustment is made for carbon-based products of incomplete combustion, such as methane, in renewable systems. This results in an under- or overestimation of the impact of CH<sub>4</sub> by 12.3% and CO by ~478% in renewable systems. This error is propagated both in scientific studies and in carbon accounting policies. We advocate first for full-carbon accounting of biomass-derived emissions, but also provide adjusted global warming impacts for emissions from proven renewable systems.

### 1 Addressing the assumptions of “carbon neutrality”

Current research on and methodologies for biofuel-based carbon accounting perpetuate a systematic error: the misapplication of the concept of “carbon neutrality.” While the fact that this concept has been applied inappropriately has been identified (Searchinger et al. 2009), we have noticed an important new error being perpetuated when accounting for the carbon (C)-based non-CO<sub>2</sub> greenhouse gas (GHG) emissions of renewable biomass fuel systems. The crux of the error is that in truly renewable biomass systems, the emission of C-based products of incomplete combustion (PICs), such as methane (CH<sub>4</sub>), would result in a lower net GHG impact than under non-renewable conditions (DeLuchi 1991; Varshney and Attri 1999), a fact which is consistently overlooked in C accounting methodologies, resulting in the

---

**Electronic supplementary material** The online version of this article (doi:10.1007/s10584-010-9984-5) contains supplementary material, which is available to authorized users.

---

T. L. Whitman (✉) · C. J. Lehmann  
College of Agriculture and Life Sciences, Cornell University, Ithaca, NY 14853, USA  
e-mail: TLW59@cornell.edu

under- and overestimation of GHG reductions in systems that rely on renewable biomass for fuel.

At present, the carbon neutral status of biofuels is often justified by the assumption that  $\text{CO}_2$  emitted from the combustion of biomass fuels was recently removed from the atmospheric  $\text{CO}_2$  stocks by plants fixing C, and thus should not be counted as a net GHG emission (e.g., Roedel 2010). This assumption can be problematic for many reasons (Friedland and Gillingham 2010; Johnson 2009; Pingoud et al. 2010; Searchinger et al. 2009). One of the foremost accounting problems occurs when the removal of biomass from some systems, though it may grow back eventually, actually results in lower net C storage over time than would have occurred under the baseline scenario with no harvesting of biomass (Searchinger 2010). When this is properly accounted for and avoided, we can consider the C in biomass to be truly “additional” (Searchinger 2010). Other sources may use the term “renewable” for this biomass C. We will use this term, because it is consistent with the literature in which the error discussed in this paper is most striking—biomass cook stove literature—but we recognize that “additional” may be a more appropriate term. (We discuss the conditions for the application of this term in more depth in Section 3.)

Carbon accounting for non- $\text{CO}_2$  GHGs has also been problematic. When combustion of fuels is not completely efficient, it produces some carbon-based PICs besides  $\text{CO}_2$ , such as  $\text{CH}_4$ , which can have higher global warming impacts than  $\text{CO}_2$ . While the carbon neutral biofuel paradigm does not count  $\text{CO}_2$  from renewable biofuel sources, in some cases, these non- $\text{CO}_2$  gases are counted directly as net GHG emissions (Caserini et al. 2010; Cherubini 2010; Gold Standard 2010; IPCC 2006; UNFCCC 2007). While this is relatively sensible for non-C-based GHGs, such as  $\text{N}_2\text{O}$ , and is an improvement over ignoring all GHG emissions from “renewable” biomass, for C-based GHGs, it is an issue, as we discuss below.

## 2 The consequences of ignoring C stoichiometry


The best way to understand this error is by considering a system in which completely renewable biomass is combusted. Under perfectly efficient combustion (Supplementary Fig. 1A), each  $\text{CO}_2$  molecule released through burning the biomass was originally fixed from atmospheric  $\text{CO}_2$  through photosynthesis and will be taken up again as the biomass re-grows, resulting in zero net emissions. The net change in the  $\text{CO}_2$  stock in the atmosphere is 0. Under the alternate scenario (Supplementary Fig. 1B), combustion is not perfectly efficient, and some of the C from the  $\text{CO}_2$  that was fixed is released upon combustion not as  $\text{CO}_2$ , but as  $\text{CH}_4$ . This  $\text{CH}_4$  has a warming effect equivalent to 25 times an equivalent mass of  $\text{CO}_2$ , based on the IPCC's 100-year global warming potential (GWP) (Forster et al. 2007). On a molar basis, rather than the standard mass basis, the warming effect of  $\text{CH}_4$  is equivalent to 9.1 times that of  $\text{CO}_2$ . Under the renewable biomass C neutrality paradigm, we would be inclined to suggest that the net impact of this system is 9.1 mol  $\text{CO}_2$ -equivalents ( $\text{CO}_2\text{e}$ ), because we ignore  $\text{CO}_2$  emissions but count all non- $\text{CO}_2$  GHG emissions in a renewable system. However, because  $\text{CH}_4$  is a C-based PIC, the emission of one molecule of  $\text{CH}_4$  means that one atom of C will not be released as  $\text{CO}_2$ . If the system is, indeed, renewable, as we have stipulated, then a  $\text{CO}_2$  molecule will still be fixed to replace the atom of C released from the plant. Thus, the net impact,

or “renewable GWP” (rGWP) of this system is actually  $9.1 - 1 = 8.1$  mol CO<sub>2</sub>e. Ignoring the biogeochemistry of the system by thinking of “methane gas” and not remembering that this stands for the C-containing chemical compound CH<sub>4</sub>, the stoichiometry of the system is lost, and its GHG impact will be overestimated.

The magnitude of this error depends heavily on the GWP of the gases under consideration. Although methane is the only C-based non-CO<sub>2</sub> GHG currently counted under the Kyoto Protocol that would be produced through combustion (HFCs and PFCs contain carbon and are counted, but would not be significant PICs), other PICs are known to have warming effects, such as non-methane hydrocarbons or carbon monoxide (CO), which enhances ozone and reduces OH levels, leading to greater CH<sub>4</sub> concentrations (Forster et al. 2007). For compounds with high GWPs, the overestimation makes only a small difference, but for compounds with low GWPs, it results in a very large error (Fig. 1). For example, in systems where all biomass burning is renewable, the estimated total GHG impact for CH<sub>4</sub> without the renewability correction factor would be 12.3% greater than its true value. If the warming impact of CO, which is estimated to lie around 1.9 by mass (Forster et al. 2007), or 1.2 on a molar basis, is considered, the estimated total GHG impact for CO from renewable sources without the renewability correction factor would be ~478% greater than its true value.

The degree to which this error affects the final estimation of a system’s impact scales directly with the fraction of the total system emissions made up by non-CO<sub>2</sub> C-based GHG emissions from renewable biomass. In very clean-burning industrial systems, CH<sub>4</sub> emissions are often relatively low compared to CO<sub>2</sub>—the IPCC default CH<sub>4</sub> emission factor for stationary combustion of solid biomass is 0.03% that of CO<sub>2</sub> by mass (IPCC 2006). However, for biomass cook stoves, this ratio may be two to three orders of magnitude higher (Johnson et al. 2008; Smith et al. 1993). In the system studied by Johnson et al. (2008), where 80% of biomass gathering is renewable, the GHG emission reductions achieved through the introduction of improved cook stoves are around 30% higher than they would be if applying the rGWP correction factor. While other uncertainties around the impact of cook stove emissions may be substantial in comparison to this error—for example, the impact

**Fig. 1** Percent overestimation of global warming impact vs. molar GWP of single-C GHGs



of particulate black carbon on climate change is poorly understood (Bond and Sun 2005)—this is no excuse to continue to use improper accounting procedures.

It is important to recognize that this error can result in an underestimation or an overestimation of GHG impact, depending on the project scenario and the baseline scenario. Overestimation occurs whenever the project scenario provides a reduction of non-CO<sub>2</sub> C-based GHG emissions from renewable biomass sources from that which was being used under the baseline scenario. An example would be if a cook stove using renewable biomass were improved to require less fuel. However, if the project scenario represents a switch to renewable biomass sources from non-renewable biomass or fossil fuel sources, then this error would cause its potential reductions to be underestimated. For example, if the baseline scenario were fossil fuel burning (all fuel non-renewable) and the project scenario were renewable biomass combustion, then counting uncorrected CH<sub>4</sub> emissions from the biomass combustion would maintain a conservative estimate of emission reductions. If the goal were simply to measure the total emissions of a system, then counting uncorrected CH<sub>4</sub> emissions from the biomass combustion would also result in a higher estimate of total emissions, which would be conservative. In general, an underestimation will result if net emissions from renewable biomass sources increase under the project scenario (although total project emissions may decrease), while a net decrease in emissions from renewable biomass sources results in an overestimation.

### 3 Righting two wrongs: addressing two carbon accounting issues

Whether this error in accounting for CH<sub>4</sub> and similar PICs is problematic or not depends partially on how CO<sub>2</sub> emissions are accounted for in the first place. We consider here three dominant approaches: (i) CO<sub>2</sub> emissions from biomass fuels are given a GHG impact value of 0, (ii) only a portion of CO<sub>2</sub> emissions from biomass fuels are counted, dependent on what portion of the biomass is from non-renewable sources, (iii) all CO<sub>2</sub> emissions are counted and all changes in biomass C storage over time are also counted. We will not discuss the specific merits or drawbacks of each approach in detail here, but, rather, focus on how the treatment of CO<sub>2</sub> emissions affects the PIC accounting error (as summarized in Table 1).

In the case of the first approach, if it is deemed justifiable to ignore emissions of CO<sub>2</sub>, and if CH<sub>4</sub> is considered to be a significant source of emissions (it may not be counted at all (Roedel 2010)), then the same justification for ignoring CO<sub>2</sub> emissions should be applied for using the rGWP or a similar accounting approach for CH<sub>4</sub> emissions. This approach has been used by some (DeLuchi 1991; Sander and Murthy 2010) but is still not applied by others (Caserini et al. 2010; Cherubini 2010; UNFCCC 2007; Gadde et al. 2009). However, as discussed in Section 2, depending on the baseline scenario, this can result in conservative emission reductions (e.g., UNFCCC 2007), in which case it is less critical that the error be righted.

The second approach begins to recognize that in many systems, particularly for wood fuel gathering, biomass comes from different sources with different degrees of renewability. This approach is currently being used in small-scale biomass fuel applications, particularly cook stove research (Johnson et al. 2009; MacCarty et al. 2008), the Kyoto Protocol's Clean Development Mechanism (CDM)'s small-scale methodology (UNFCCC 2009) and the Gold Standard's carbon offset methodology

**Table 1** Accounting approaches for CO<sub>2</sub> emissions and C-based non-CO<sub>2</sub> GHG emissions (e.g., CH<sub>4</sub>) in biomass fuel systems

Current accounting for CO <sub>2</sub> emissions	Current and recommended accounting for non-CO <sub>2</sub> C-based GHG emissions
CO <sub>2</sub> emissions are given a value of 0 because biomass stocks are determined to be “renewable” or “additional”.	C-based non-CO <sub>2</sub> GHG emissions are not included (Roedl 2010) or are included and weighted either using the full GWP (Caserini et al. 2010; Cherubini 2010; Gadde et al. 2009) or the rGWP (DeLuchi 1991; Sander and Murthy 2010). If PIC impacts are included (and CO <sub>2</sub> neutrality is justified), rGWP should be applied, particularly where this results in a conservative estimate of emission reductions.
CO <sub>2</sub> emissions are partly counted (non-renewable biomass) and partly valued at 0 (renewable biomass), depending on fNRB of biomass fuel.	C-based non-CO <sub>2</sub> GHG emissions from non-renewable biomass are included in all cases. Emissions from renewable biomass are excluded (UNFCCC 2009) or included with full GWP weighting (MacCarty et al. 2008; Johnson et al. 2009; Gold Standard 2010) or included with a reduced GWP (Whitman 2010). The impact of PIC emissions from renewable biomass should be weighted using the rGWP, particularly where this results in a conservative estimate of emission reductions.
All C flows within the system are modeled explicitly over time.	The issue of miscounting non-CO <sub>2</sub> C-based GHGs is avoided by modeling the C fixation by plants directly (Bailis 2009; González-García et al. 2010).

for improved cook stoves (Gold Standard 2010). It combines the carbon neutrality paradigm with estimates of the degree of renewability of the biomass used for fuel (Johnson et al. 2010a).

A measure of the fraction of fuel that is from “non-renewable” biomass (fNRB) is used to determine the fraction of CO<sub>2</sub> emissions that should be counted as net emissions, while remaining, “renewable” CO<sub>2</sub> emissions are ignored. In order for this principle to function appropriately, if any biomass designated as “renewable” is gathered from a stock in perpetuity, the stock should both be replenished, and also not have increased beyond its stable level if the gathering were not to have taken place (i.e., be truly “additional” biomass (Searchinger 2010)). If biomass designated as “non-renewable” is gathered from a stock, it should immediately deplete the stock, and the stock should never be replenished. However, these extreme scenarios do not usually apply neatly to a given region, so certain conventions are used to estimate fNRB.

Projects under the CDM consider NRB to be biomass that is not demonstrably renewable and meets at least two out of three other conditions. Demonstrably renewable biomass is defined for forests as cases where: (i) the land area remains a forest, (ii) sustainable management ensures C stocks do not decrease over time,

although they may temporarily decrease due to harvesting, (iii) locally applicable forestry and nature conservation regulations are complied with. In addition to not being demonstrably renewable biomass, the three conditions to designate non-renewable biomass are: (i) users spend increasing time or travel increasing distances to gather or transport wood into the project area, (ii) data show that C stocks are being depleted in the project area, (iii) fuel wood price is increasing (UNFCCC 2009).

In these cases where biomass is classified as being renewable or non-renewable, non-CO<sub>2</sub> GHG emissions are not treated consistently. The CDM cook stove approach does not consider any emissions from renewable biomass (UNFCCC 2009), while others consider the non-CO<sub>2</sub> emissions from both renewable and non-renewable biomass sources (Gold Standard 2010; Johnson et al. 2009; MacCarty et al. 2008; Whitman 2010). For non-renewable fuel emissions, CH<sub>4</sub> emissions are weighted by their full GWP and there is no error. However, when renewable fuel emissions are counted, CH<sub>4</sub> emissions should be valued according to the rGWP wherever CO<sub>2</sub> emissions are ignored. The CDM's current methodology remains conservative, because it does not include emissions from any renewable biomass (UNFCCC 2009), but the Gold Standard cook stove methodology and other studies currently overestimate any emission reductions from renewable biomass sources (Gold Standard 2010; Johnson et al. 2008; MacCarty et al. 2008).

Both these first two approaches rely on assumptions about the renewability status of the system. Perhaps the most robust way of addressing such an error is the third approach—to apply full carbon accounting techniques, where the net flows of carbon are explicitly traced over time, including the CO<sub>2</sub> and non-CO<sub>2</sub> emissions from using bioenergy as well as the increases or decreases in net fixation of CO<sub>2</sub> by biological carbon stock due to land-use changes (González-García et al. 2010; Guinée et al. 2009; Johnson 2009; Searchinger et al. 2009). If a biofuel is truly carbon-neutral, then this fact will emanate from the C balance sheet and there is no need to treat CH<sub>4</sub> emissions differently. This system-based solution is an ideal approach for scientific certainty, and can be applied at the project scale by measuring or modelling biomass growth and fuel emissions (Bailis 2009).

#### 4 Fixing the carbon accounting error

Substantially different approaches are being used in order to determine the C budget of renewable biomass fuel systems, with varying consequences. As discussed above, the optimal approach to avoid many C accounting errors and increase confidence in the C accounting for offsets or national inventories would be to use full-system carbon accounting, by explicitly modeling or measuring biomass and soil C growth, decay, and loss as well as emissions.

If renewability status or the fNRB can be justified and are used, however, the proposed rGWP corrections must be applied in place of GWPs for PICs from renewable biomass in order to correct this error (Table 2). (See Shine (2009) for a critique of GWPs and Levasseur et al. (2010) for a discussion of applying GWPs to measure the impacts of gases emitted over time.) To avoid underestimating PIC emission impacts, the rGWP should only be used where biomass can be explicitly proven to be renewable or where applying the correction results in a conservative estimate of emission reductions, such as in improved biomass cook stove systems.

**Table 2** One hundred-year global warming potential (GWP) (Forster et al. 2007) and renewable biomass global warming potential (rGWP) on a mass and molar basis

Compound	GWP (t CO <sub>2</sub> e/t GHG)	rGWP (t CO <sub>2</sub> e/t GHG)	GWP (mol CO <sub>2</sub> e/mol GHG)	rGWP (mol CO <sub>2</sub> e/mol GHG)
CH <sub>4</sub>	25	22.26	9.12	8.12
CO	1.9	0.33	1.21	0.21

This recommendation should be added to those discussed by Johnson et al. (2010a, b) for revising the current accounting procedures for biomass cook stove systems, and to those recommended by Searchinger et al. (2009) for improving GHG inventories.

**Acknowledgements** This research was supported by a Natural Sciences and Engineering Research Council of Canada PGS-M grant and by the Cornell Center for a Sustainable Future (T.W.). Thanks to two anonymous reviewers for their helpful comments and suggestions.

## References

- Bailis R (2009) Modeling climate change mitigation from alternative methods of charcoal production in Kenya. *Biomass Bioenergy* 33:1491–1502
- Bond T, Sun H (2005) Can reducing black carbon emissions counteract global warming? *Environ Sci Technol* 39:5921–5926
- Caserini S, Livio S, Giugliano M, Grosso M, Rigamonti L (2010) LCA of domestic and centralized biomass combustion: the case of Lombardy (Italy). *Biomass Bioenergy* 34:474–482
- Cherubini F (2010) GHG balances of bioenergy systems—overview of key steps in the production chain and methodological concerns. *Renew Energy* 35:1565–1573
- DeLuchi MA (1991) Emissions of greenhouse gases from the use of transportation fuels and electricity, vol 1: summary. Argonne National Laboratory, Argonne, pp 77–78
- Forster PV, Ramaswamy V, Artaxo P, Berntsen T, Betts R, Fahey DW, Haywood J, Lean J, Lowe DC, Myhre G, Nganga J, Prinn R, Raga G, Schulz M, Van Dorland R (2007) Changes in atmospheric constituents and in radiative forcing. In: Solomon SD et al (eds) *Climate change 2007: the physical science basis*. Contribution of working group I to the fourth assessment report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, pp 129–234
- Friedland AJ, Gillingham KT (2010) Carbon accounting a tricky business. *Science* 326:411
- Gadde G, Menke C, Wassman R (2009) Rice straw as a renewable energy source in India, Thailand, and the Philippines: overall potential and limitations for energy contribution and greenhouse gas mitigation. *Biomass Bioenergy* 33:1532–1546
- Gold Standard (2010) *Methodology for improved cook-stoves and kitchen regimes, vol 2*. Geneva, Switzerland
- González-García S, Luo L, Moreira MR, Feijoo G, Huppes G (2010) Life cycle assessment of flax shives derived second generation ethanol fueled automobiles in Spain. *Renew Sustain Energy Rev* 13:1922–1933
- Guinée JB, Heijungs R, van der Voet E (2009) A greenhouse gas indicator for bioenergy: some theoretical issues with practical implications. *Int J Life Cycle Assess* 14:328–339
- IPCC (2006) *IPCC guidelines for national greenhouse gas inventories*. National Greenhouse Gas Inventories Programme, Institute for Global Environmental Strategies, Tokyo, Japan
- Johnson E (2009) Goodbye to carbon neutral: getting biomass footprints right. *Environ Impact Assess Rev* 29:165–168
- Johnson M, Edwards R, Frenk CA, Masera O (2008) In-field greenhouse gas emissions from cook-stoves in rural Mexican households. *Atmos Environ* 42:1206–1222
- Johnson M, Edwards R, Ghilardi A, Berrueta V, Gillen D, Frenk CA, Masera O (2009) Quantification of carbon savings from improved biomass cookstove projects. *Environ Sci Technol* 43:2456–2462

- Johnson M, Edwards R, Masera O (2010a) Improved stove programs need robust methods to estimate carbon offsets. *Clim Change*. doi:10.1007/s10584-009-9770-4
- Johnson M, Edwards R, Berrueta V, Masera O (2010b) New approaches to performance testing of improved cookstoves. *Environ Sci Technol* 44:368–374
- Levasseur A, Lesage P, Margni M, Deschênes L, Samson R (2010) Considering time in LCA: dynamic LCA and its application to global warming impact assessments. *Environ Sci Technol* 44:3169–3174
- MacCarty N, Ogle D, Still D (2008) A laboratory comparison of the global warming impact of five major types of biomass cooking stoves. *Energy Sustain Dev* 12:5–14
- Pingoud K et al (2010) Bioenergy: counting on incentives. *Science* 327:1199
- Roedel A (2010) Production and energetic utilization of wood from short rotation coppice—a life cycle assessment. *Int J Life Cycle Assess* 15:567–578
- Sander K, Murthy GS (2010) Life cycle analysis of algae biodiesel. *Int J Life Cycle Assess* 15:704–714
- Searchinger TD (2010) Biofuels and the need for additional carbon. *Environ Res Lett* 5:024007 (10pp). doi:10.1088/1748-9326/5/2/024007
- Searchinger T, Hamburg SP, Melillo J, Chameides W, Havlik P, Kammen DM, Likens GE, Lubowski RN, Obersteiner M, Oppenheimer M, Robertson GP, Schlesinger WH, Tilman GD (2009) Fixing a critical climate accounting error. *Science* 326:527–528
- Shine KP (2009) The global warming potential—the need for an interdisciplinary retrieval. *Clim Change* 96:467–472
- Smith KR, Khalil MAK, Rasmussen RA, Thorneloe SA, Mandgdeg F, Apte M (1993) Greenhouse gases from biomass and fossil fuel stoves in developing countries: a Manila plot study. *Chemosphere* 26:479–505
- UNFCCC (2007) Grid-connected electricity generation using biomass from newly developed dedicated plantations. AM0042/version 02, sectoral scopes: 01 & 14, Clean Development Mechanism Executive Board 35
- UNFCCC (2009) Energy efficiency measures in thermal applications of non-renewable biomass. ILG./version 02, sectoral scope: 03, Clean Development Mechanism Executive Board 51
- Varshney CK, Attri AK (1999) Global warming potential of biogenic methane. *Tellus B* 51:612–615
- Whitman TW (2010) Biochar as a carbon sequestration mechanism: decomposition, modelling, and policy. Unpublished master's thesis, Cornell University, Ithaca